

## PATIENT INFORMATION LEAFLET

### SELOPARIN 25 000 IU/5 ml solution for injection.

#### Sterile

#### It is for subcutaneous and intravenous administration.

Each 5 ml vial contains:

- ❖ **Active substance:** 25000 IU heparin sodium
- ❖ **Excipients:** Benzyl alcohol, sodium chloride, water for injections and sodium hydroxide or hydrochloric acid to keep acidity on normal levels.

**Read all of this SELOPARIN carefully before you are given this medicine, because it contains important information for you.**

- *Keep this leaflet. You may need to read it again.*
- *If you have any further questions, please ask your doctor or pharmacist.*
- *This medicine has been prescribed for personally; you should not pass it on to others.*
- *During use of this medicine please tell to your doctor that you use this medicine when you go to the doctor or hospital.*
- *Follow the information in the leaflet exactly. Do not use **lower or higher** dosages regarding the medicine.*

#### **In this Patient Information Leaflet:**

- 1. *What SELOPARIN is and what is it used for?***
- 2. *What you need to know before you use SELOPARIN?***
- 3. *How to use SELOPARIN?***
- 4. *What are the possible side effects?***
- 5. *How to store SELOPARIN?***

**Headings are provided.**

#### **1. What SELOPARIN is and what is it used for?**

SELOPARIN belongs to a group of drugs that are called anticoagulants as heparin sodium. SELOPARIN vial contains heparin sodium in a liquid solution. The box contains one vial. SELOPARIN is standard heparin.

Heparin sodium helps to stop blood clotting. Blood flow is allowed to continue uneventfully.

Blood carrying veins in body are named arteries or veins.

SELOPARIN is used in the following cases:

- To prevent further development of harmful blood clots in veins (arteries or veins) (for treatment)
- To help to stop harmful blood clots in veins (arteries or veins) (preventive)
- Hemodialysis and heart surgery
- If the risk of heart attack or stroke are available
- Several blood coagulation disorders

Blood clots in the body can occur in leg veins (deep vein thrombosis, DVT), in the lungs (pulmonary embolism).

If you have more weight or are pregnant or have some blood diseases or have had DVT or pulmonary embolism. Heart attack, paralysis, long term inactivity can increase clot risk.

## 2. What you need to know before you use SELOPARIN?

*If you are under epidural and spinal anesthesia;*

*Please tell your doctor, If you are pregnant and/or you have used heparin before taking an anaesthetic.*

*Please inform to person who gives narcosis (dentist included) that you received SELOPARIN in case of planned operation.*

Health personnel will make regular checks after taken anesthesia. It will be checked whether purpleness and bleeding around spine.

These problems can cause irreversible paralysis. This situation can occur like tremble (shiver) on legs or body, weakness or numbness, back pain or urinary problem. But they are seen very rarely.

Your doctor will tell you when you can take medicine after anesthesia

### **YOU SHOULD NOT BE GIVEN SELOPARIN if you:**

- have hypersensitivity to heparin sodium or any of the ingredients of SELOPARIN (Benzyl alcohol)
- have or have ever had thrombocytopenia (a serious blood disorder which prevents blood from clotting properly)
- have hemophilia (a genetic disorder which may cause you to bleed excessively)

- have very high blood pressure
- have serious liver problems
- have peptic ulcer
- have known endocarditic (inflammation of the covering layer of inner surface of the heart and heart valves)
- have ever had bleeding into the brain.
- have injuries in the spine, brain, eyes and ears
- have recently had surgery of the brain, spine eyes or ears or you will have surgery like this.
- have the possibility of miscarriage.

Heparin sodium cannot be given to premature, newborn babies or babies of up to one month.

**TAKE SPECIAL CARE with SELOPARIN if you:**

- suffer from allergies or have previously had an allergic reaction to low molecular weight (LMW) heparin such as Tinzaparin, enoksaparin or deltapirin
- have kidney problems
- have liver problems
- have eye disorders (Retina)
- have blood pressure problems (high pressure)
- suffer from diabetes
- have recently had lumbar puncture
- have metabolic acidosis
- suffer from high levels of potassium in your blood or are taking a medicine that may increase the potassium level in your blood (e.g. amiloride, spironolactone).
- have any condition which makes you to bleed more easily
- will use for 1 to 3 years old children, please tell your doctor
- are taking another medicines affecting blood coagulation at the same time with heparin, please check ‘Application while taking other medicines’ for these medicines list.
- are pregnant or wish to become pregnant.

Please read ‘important informations about excipients contained in SELOPARIN’

Your doctor will do your blood tests before application of Seloparin and during your use. Your doctor can check medicine’s dose. These tests check coagulation cells (thrombocyte) and potassium level of the blood.

This medicine can cause to bleed easily. Your doctor or your nurse should be careful when other injection or operation are applied.

If these warnings are available for any period in the past. Please tell your doctor.

**Administration of SELOPARIN with food and beverages**

Heparin cannot be given if you drink large amounts of alcohol.

## **Pregnancy**

*Please consult your doctor or pharmacist before using this medicine.*

Your doctor can apply heparin during pregnancy, if necessary is. After using heparin, epidural anesthesia especially cannot be applied during birth.

*Please consult your doctor or pharmacist if you notice that you are pregnant during treatment.*

## **Breast-feeding**

*Please consult your doctor or pharmacist before using this medicine.*

It can be used during breastfeeding.

## **Driving and using machines**

SELOPARIN usually can have low effect on driving or using machines. Please consult your doctor if you feel a side effect which retains using car.

## **Important information about some of the excipients in SELOPARIN**

This medicinal product contains 33,21 mg sodium in each 5 ml vial. This situation should be considered if patients are on sodium diet.

SELOPARIN contains benzyl alcohol (9,45mg/ml) as preservative. Please be careful when suspected patients are prescribed for. It cannot be applied for premature and newborn babies.

It may cause allergic reactions and toxic reactions for babies and children up to 3 years old.

If you are pregnant, you doctor will decide to apply Seloparin.

## **Administration while taking other medicines**

- If you use angiotensin converting enzyme inhibitor or enalapril, losartan or valsartan called angiotensin II antagonists for high blood pressure or heart problems treatment, your potassium blood level can increase.(hyperkalemia)
- If you use ibuprofen or diclofenac called non-steroidal anti-inflammatory drugs for arthritis or other pains, possibility of bleeding can be very easily.
- If you use aspirin called drugs of salicylate group to prevent blood clot or to reduce pain or inflammation, possibility of bleeding can be very easily.
- If you use clopidogrel called platelet aggregation inhibitors to stop harmful blood clot, possibility of bleeding can be very easily.
- If you use streptokinase called thrombolytic agents to lyse blood clot, possibility of bleeding is very high level.
- If you use warfarin called vitamin K antagonist to stop harmful blood clot, possibility of bleeding is very high level.
- Glyceryl trinitrate infusion for angina treatment can increase heparin's effect.
- Activated protein C: to get rid of blood clot. Possibility of bleeding can be very easily.
- If you take dextran to increase blood volume, possibility of bleeding can be very easily.

- Corticosteroids (gluco-): excluding hydrocortisone used for replacement therapy in Addison's disease.
- Cephalosporins: Some cephalosporins, e.g. cefaclor, cefixime and ceftriaxone, can affect the coagulation process and may therefore increase the risk of hemorrhage when used concurrently with heparin.
- Tobacco smoke: Nicotine may partially counteract the anticoagulant effect of heparin. Increased heparin dosage may be required in smokers.

If you take above medicines at the same time as heparin, your doctor can do check up tests included blood tests.

*Please tell your doctor or pharmacist if you are taking, or have recently taken, any other medicines including medicines obtained with or without prescription.*

### **3. How to use SELOPARIN?**

#### **Instructions for the appropriate usage and dose / frequency of application:**

Heparin concentration in SELOPARIN is 5 000 IU/ml. All heparin preparations do not contain same concentration. Because of that heparin dose is stated as IU.

SELOPARIN will be applied from doctor or nurse.

Your doctor will decide the dose of medicine given to you depending on your medical condition and will apply to you.

SELOPARIN cannot be injected mixing with another medicines.

#### **Administration route and method of administration:**

SELOPARIN is applied by intravenously or subcutaneous.

#### **Various age groups:**

##### **Children**

SELOPARIN cannot be used for children under one month.

##### **The elderly**

For elderly patients, the dose should be reduced. aPTT (activated partial thromboplastin time) should be followed.

#### **Administration for certain groups**

##### **Renal and Liver failure**

If you have kidney or liver problems. Your doctor can decide to be taken lower dose SELOPARIN.

*Please talk to your doctor or pharmacist if you think that the effect of SELOPARIN is too strong*

*or too weak.*

### **If you think you have been given too much SELOPARIN**

Your doctor or nurse do not apply SELOPARIN at high doses.

If you think that SELOPARIN is given to you at high doses. Please talk your doctor or your nurse.

Hemorrhage (serious bleeding) may occur. (Please look 4.section)

Protamine sulphate can be applied with another injection.

*Please talk to your doctor and pharmacist if you use SELOPARIN more than you need to use.*

### **If you forget to take SELOPARIN**

Your doctor or nurse do not apply SELOPARIN at high doses.

If you forget to take a dose. Please talk your doctor or nurse.

*Please talk to your doctor and pharmacist if you have any questions for implementation of this medicine*

## **4. What are possible side effects?**

Like all medicines, side effects can be seen in people who are sensitive to the ingredients of SELOPARIN.

Please inform your doctor or pharmacist if you have any side effect, worsening of one of the side effects or any side effect that is not mentioned in this leaflet.

**If any of the following cases occurs, discontinue SELOPARIN, IMMEDIATELY inform your physician or admit to the emergency service of the closest hospital:**

- Difficulty breathing
- Swelling of face and throat
- Severe skin rash
- Bubbles around the injection site.

All of these are very serious side effects.

If you have one of these, it means you have a serious allergy against SELOPARIN. You may need urgent medical attention or hospitalization.

**Tell your doctor immediately or refer to the emergency of nearest hospital if you notice any of the following after epidural anesthesia or spinal anesthesia. Paralysis may occur:**

- Tingling ,weakness or numbness (legs or down from the waist)
- Back pain
- Urinary problems.

**Tell your doctor if you notice any of followings. It means serious bleeding:**

- Red or brown urine
- Black feces like tar
- Unusual bruising
- Unceasing bleeding in the nose and mouth or surgical wound

**Other side effects include:**

**Very common side effects (affects 1 user in 10 patients)**

- Injection site reactions: Irritation may occur when injected locally under the skin.

**Common side effects (less than 1 user in 10 patients but more than 1 user in 100 patients)**

- Rash
- Itchy raised rashes (urticaria)
- The reduction of bone density (osteoporosis). Bones of patients treated with heparin for a long time can be broken easily and can be occurred power loss.

**Uncommon side effects (less than 1 user in 100 patients but more than 1 user in 1000 patients)**

- Decreased bone density (osteoporosis). Patients who use heparin for a long time will lose their bones and break easily.
- Hair loss

**Rare side effects (less than 1 user in 10000 patients)**

- Easy bruising or easy bleeding. Harmful blood clot may occur. Cells providing the coagulation (thrombocytes) decrease. Your doctor can explain this situation.
- Changes in the blood test results. Potassium level can increase. If you have serious kidney problems or diabetes, change possibility of the blood test results may be more high. Your doctor can explain this situation.
- Various hypersensitivity symptoms and severe allergic reactions (conjunctivitis, rhinitis, asthma, tachycardia, fever, etc.), anaphylactoid reactions and anaphylactic shock (very serious allergic reactions)
- Increased serum transaminase, gamma-GT, LDH and lipase values (some enzymes that can be measured by blood tests) may occur; but usually resolve on discontinuation of heparin. If you are unsure, ask your doctor.
- Bleeding can occur. Especially too much heparin can cause bleeding.
- Necrosis of skin may occur. If this occurs treatment should be discontinued immediately.

**Very rare (< 1/1000 but ≥1/10.000)**

- Long time painful erections in men

- Anaphylactoid reactions and anaphylactic shock (very severe allergic reactions)

*Please talk to your doctor if you think that any of the above side effects may occur. Some side effects can be serious.*

*Please inform your doctor or pharmacist, if you notice any side effects not listed in this leaflet.*

### **Reporting of the side effects**

If you get any side effects not listed in this leaflet, talk to your doctor or pharmacist. You can also report side effects directly to your doctor or pharmacist. You can also report side effects directly to your country's related health authority. By reporting side effects, you can help provide more information on the safety of this medicine

### **5. How to store SELOPARIN?**

*Keep SELOPARIN out of the reach and sight of children. Keep SELOPARIN in the package.*

Keep at room temperature below 25° C. Protect from freezing.

Chemical and physical in use stability has been demonstrated for 28 days at 25°C

Please be careful that solution is clear and has unopened packaging before application.

Multidose vial is for use of one patient.

### **Use in accordance with expiration dates.**

*Do not use after the expiry date on the packaging of SELOPARIN. Do not use SELOPARIN if you notice any damage at the packaging.*

### ***Marketing Authorization Holder:***

Haver Farma İlaç A.Ş.  
Akbaba Mah. Maraş Cad. No:52/2/1  
Beykoz / ISTANBUL / TURKEY

### ***Manufacturing Site:***

Mefar İlaç San.A.Ş.  
Ramazanoğlu Mah. Ensar Cad. No:20,  
Kurtköy-Pendik / ISTANBUL / TURKEY

*This patient information leaflet is approved at 19/01/2016.*





## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

SELOPARIN 25 000 IU/5 ml solution for injection

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ampoule (5 ml) contains;

#### Active ingredient:

25 000 IU heparin sodium

#### Excipients:

Benzyl alcohol                      47.25 mg

Sodium chloride                      45 mg

For a full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Solution for 5 ml ampoule

Almost colourless, clear solution for injection

### 4. CLINICAL PARTICULARS

#### 4.1. Therapeutic indications

SELOPARIN is indicated mainly at diseases below:

- Prophylaxis of deep vein thrombosis, pulmonary embolism and arterial thromboembolic events.
- Treatment of deep vein thrombosis, pulmonary embolism, unstable angina pectoris and acute peripheral arterial occlusion.
- Prophylaxis of mural thrombosis following myocardial infarction.
- In extracorporeal circulation and haemodialysis.

#### 4.2. Posology and method of administration

##### Posology:

Use intravenously or subcutaneously. It should not be administered intramuscularly.

The concentration of heparin solution is 5000 IU / ml. Heparin dose as IU must be specified for all heparin preparations does not contain same concentration.

**Application frequency and time:**

- Prophylaxis of deep vein thrombosis, pulmonary embolism and arterial thromboembolic events:

2 hours preoperatively: 5,000 units subcutaneously

followed by: 5,000 units subcutaneously every 8-12 hours, for 7-10 days or until the patient is fully ambulant.

No laboratory monitoring should be necessary during low dose heparin prophylaxis. If monitoring is considered desirable, anti-Xa assays should be used as the activated partial thromboplastin time (APTT) is not significantly prolonged.

Laboratory tests should be performed daily. Ideally, every day the same hours and 6 hours after the start of the first sample treatment, then after each dose in exchange for 4-6 hours should be taken. During the whole dose of heparin therapy, daily laboratory monitoring, activated part thromboplastin time (aPTT) adjusting or controlling 1.5-2 times the value of the midpoint of the normal range is essential.

- Treatment of deep vein thrombosis, pulmonary embolism, unstable angina pectoris and acute peripheral arterial occlusion:

Loading dose: 5,000 units intravenously (10,000 units may be required in severe pulmonary embolism)

Maintenance: 1,000-2,000 units/hour by intravenous infusion,  
or 10,000-20,000 units 12 hourly subcutaneously,  
or 5,000-10,000 units 4hourly by intravenous injection.

- Prophylaxis of mural thrombosis following myocardial infarction:

12,500 units 12 hourly subcutaneously for at least 10 days.

- In extracorporeal circulation and haemodialysis:

Cardiopulmonary bypass: Initially 300 units/kg intravenously, adjusted thereafter to maintain the activated clotting time (ACT) in the range 400-500 seconds.

Haemodialysis and haemofiltration:

Initially 1,000-5,000 units,

SELOPARIN 25000 IU/ 5 ml Solution for Injection



## Module 1 Administrative Information

### Module 1.3 Product Information

#### Module 1.3.1 SPC, PIL, Ampoule Print Form, The printing on the box

Maintenance: 1,000-2,000 units/hour, adjusted to maintain clotting time >40 minutes.

#### **Route of administration:**

By continuous intravenous infusion in 5% glucose or 0.9% sodium chloride or by intermittent intravenous injection, or by subcutaneous injection.

Heparin intravenous injection volume should not exceed 15 ml.

As the effects of heparin are short lived, administration by intravenous infusion or subcutaneous injection is preferable to intermittent intravenous injections.

#### **Additional information on special populations:**

##### **Renal / Hepatic impairment**

It should be administered with caution. In patients with advanced renal or hepatic disease, a reduction in dosage may be necessary. The risk of bleeding is increased with severe renal impairment and in the elderly (particularly elderly women).

##### **Pediatric population**

- Prophylaxis of deep vein thrombosis, pulmonary embolism and arterial thromboembolic events:

No dosage recommendations.

- Treatment of deep vein thrombosis, pulmonary embolism, unstable angina pectoris and acute peripheral arterial occlusion:

Loading dose: 50 units/kg intravenously

Maintenance: 15-25 units/kg/hour by intravenous infusion,  
or 250 units/kg 12 hourly subcutaneously  
or 100 units/kg 4hourly by intravenous injection.

##### **Pregnant**

5,000-10,000 units every 12 hours, subcutaneously, adjusted according to aPTT or anti-Xa assay.

#### **4.3. Contraindications**

SELOPARIN should not be used in the following cases:

- Hipersensitivity to heparin or excipients of drug content.
- Premature babies or newborns (because it contains benzyl alcohol)

- Patients who consume large amounts of alcohol, who are sensitive to the drug, who are actively bleeding or who have haemophilia or other bleeding disorders, severe liver disease (including oesophageal varices), purpura, severe hypertension, active tuberculosis or increased capillary permeability.
- Patients with present or previous thrombocytopenia. The rare occurrence of skin necrosis in patients receiving heparin contraindicates the further use of heparin either by subcutaneous or intravenous routes because of the risk of thrombocytopenia. Because of the special hazard of postoperative haemorrhage heparin is contraindicated during surgery of the brain, spinal cord and eye, in procedures at sites where there is a risk of bleeding, in patients that have had recent surgery, and in patients undergoing lumbar puncture or regional anaesthetic block.

The relative risks and benefits of heparin should be carefully assessed in patients with a bleeding tendency or those patients with an actual or potential bleeding site eg. hiatus hernia, peptic ulcer, neoplasm, bacterial endocarditis, retinopathy, bleeding haemorrhoids, suspected intracranial haemorrhage, cerebral thrombosis or threatened abortion.

#### **4.4. Special warnings and precautions for use**

Heparin-treated (curative or prophylactic doses) in a patient;

- if ischemic stroke,
- if myocardial infarction,
- if acute ischemia in the lower limbs,
- if pulmonary embolism,
- if phlebitis,
- if thrombotic events such as thrombosis aggravation,

Heparin-induced thrombocytopenia (HIT) should be systematically considered and platelets (platelet) count test should be carried out urgently.

#### **Precautions**

*At risk situations:*

Lumbar puncture performance should be considered for intraspinal bleeding risk in postoperative period after the brain and spinal cord surgery or in patients with peptic ulcer, choroidal and retinal vascular disease a history. It should be delayed as much as possible.

*Monitoring of platelet counts:*

Regardless of the administered dose and indication, platelets depending on the risk of HIT should be monitored. Once before the treatment, then 2 times a week for 21 days, platelet counts should be performed. After this period, in which case the exact extension of the treatment, the frequency of monitoring of the platelet count should be increased to 1 per week until treatment is stopped.

Heparin can suppress adrenal secretion of aldosterone. In this case leads to hyperkalemia and/or metabolic acidosis with hypoaldosteronism. This event was observed in high risk patients with high blood levels of potassium (diabetic patients, patients with chronic renal failure, patients with metabolic acidosis story, patients using medications that increase the blood level of potassium such as ACE inhibitors and NSAID). The risk of hyperkalemia increases during treatment and is usually reversible. Plasma potassium levels should be monitored in case of prolonged treatment in high risk patients. Plasma potassium levels should be measured before the heparin treatment for patients at risk and all patients treated for more than 7 days.

*Heparin-induced thrombocytopenia (HIT):*

There is the risk of immunological origin heparin-induced thrombocytopenia, known as type II thrombocytopenia, causing thrombosis (unfractionated heparin and less frequently caused by low molecular weight heparins) sometimes.

HIT is determined by two consecutive measurements with a significant decrease in platelet count by 30-50% and/or <100 000 worth of platelet count. It develops between 5-21th days following administration of heparin (with a peak incidence of 10 days), but during the treatment with heparin in patients with a history of thrombocytopenia may occur earlier. As a result, a story of this type should be investigated systematically during in-depth interviews. In addition, the risk of relapse to drug rechallenge case can take several years or even a lifetime (see also Contraindications).

Platelet measurement must be done in 5 days more than heparin-treated patients. Treatment should be discontinued immediately in patients with thrombocytopenia development.

In all cases, the beginning of HIT is an emergency and requires specialist advice.

Any significant decrease in platelet count (initial value of 30-50%, i) should be seen as a signal before the value reaches critical levels. The observed decrease in the number of platelets;

- Emergency a platelet count
- If the decrease is confirmed, require cessation of heparin therapy.

Under these circumstances, things to emergency measures does not depend on the results of in vitro or immunological platelet aggregation tests. Because only some specialized laboratories routinely conducts these tests and the results obtained after a few years the best possibility. These tests must still be performed to aid in the diagnosis of complications. If you continue to treat, thrombosis is a major risk.

- Treatment and prevention of thrombotic complications of HIT

If deemed necessary to continue anticoagulation, heparin must be replaced with another class of antithrombotic agents. Danaparoid sodium or hirudin, depending on the situation is given in prophylactic or curative doses.

Replaced by oral anticoagulant drugs depending on the risk of thrombotic cases exacerbated with oral anticoagulants should occur only once, when platelet count returned to normal.

Although even though rare heparin to hypersensitivity, a trial dose of 1000 IU is recommended to be administered patients with a history of allergy. Care must be exercised in patients with hypersensitivity to low molecular weight heparins.

Recommended low dose regimen, the majority of patients, does not create a change in coagulation time. However, the effect on blood coagulation therapy should be kept under observation in major surgery of patients with a individual response to heparin.

Care should be taken in case of spinal or epidural anesthesia (risk of spinal hematoma).

#### *Heparin resistance*

There is considerable variation in individual anticoagulant responses to heparin.

Heparin resistance, defined as an inadequate response to heparin at a standard dose for achieving a therapeutic goal occurs in approximately 5 to 30% of patients.

Factors predisposing to the development of heparin resistance, include:

- Antithrombin III activity less than 60% of normal (antithrombin III dependent heparin resistance):

Reduced antithrombin III activity may be hereditary or more commonly, acquired (secondary to preoperative heparin therapy in the main, chronic liver disease, nephrotic syndrome, cardiopulmonary bypass, low grade disseminated intravascular coagulation or drug induced, e.g. by aprotinin, oestrogen or possibly nitroglycerin)

- Patients with normal or supranormal antithrombin III levels (antithrombin III in dependent heparin resistance)
- Thromboembolic disorders
- Increased heparin clearance
- Elevated levels of heparin binding proteins, factor VIII, von Willebrand factor, fibrinogen, platelet factor 4 or histidinerich glycoprotein
- Active infection (sepsis or endocarditis)
- Preoperative intraaortic balloon counterpulsation
- Thrombocytopenia
- Thrombocytosis
- Advanced age
- Plasma albumin concentration 35g/dl

Heparin resistance is also often encountered in acutely ill patients, in patients with malignancy and during pregnancy or the postpartum period.

SELOPARIN contains 47.25 mg benzyl alcohol as preservative. Caution needed when prescribing to suspected patients, especially infants. Benzyl alcohol causes toxic and anaphylactoid reactions in babies and < 3 years old children.

This medicinal product contains 33.21 mg sodium per 5 ml ampoule, equivalent to 0.066 % of the WHO recommended maximum daily intake of 2 g sodium for an adult.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Analgesics: Drugs that interfere with platelet aggregation eg. acetylsalicylic acid (analgesic and antipyretic doses), acetylsalicylic acid (at a dose which inhibits platelets) and other NSAIDs should be used with care. Increased risk of haemorrhage with ketorolac (avoid concomitant use even with lowdose heparin).

Corticosteroids (gluco-): excluding hydrocortisone used for replacement therapy in Addison's disease.

Anticoagulants, platelet inhibitors, etc: Increased risk of bleeding with oral anticoagulants, epoprostenol, clopidogrel, ticlopidine, streptokinase, dipyridamole, dextran solutions, or any other drug which may interfere with coagulation.

Cephalosporins: Some cephalosporins, e.g. cefaclor, cefixime and ceftriaxone, can affect the coagulation process and may therefore increase the risk of haemorrhage when used concurrently with heparin.

ACE inhibitors: Hyperkalaemia may occur with concomitant use.

Nitrates: Reduced activity of heparin has been reported with simultaneous intravenous glyceryl trinitrate infusion.

Thrombolytic drugs: It may increase the risk of hemorrhagic. Patients should be monitored regularly.

Probenecid: May increase the anticoagulant effects of heparin.

Tobacco smoke: Nicotine may partially counteract the anticoagulant effect of heparin. Increased heparin dosage may be required in smokers.

Interference with diagnostic tests may be associated with pseudohypocalcaemia (in haemodialysis patients), artefactual increases in total thyroxine and triiodothyronine, simulated metabolic acidosis and inhibition of the chromogenic lysate assay for endotoxin. Heparin may interfere with the determination of aminoglycosides by immunoassays.

#### **4.6. Pregnancy and lactation**

##### **General advice**

Pregnancy category: C

##### **Women of childbearing potential / Birth control (Contraception)**

There is no study concerning the effect of heparin on women with childbearing potential/contraception.

##### **Pregnancy**



Heparin is not contraindicated in pregnancy. Heparin does not cross the placenta. The decision to use heparin in pregnancy should be taken after evaluation of the risk/benefit in any particular circumstances.

Reduced bone density has been reported with prolonged heparin treatment during pregnancy.

Haemorrhage may be a problem during pregnancy or after delivery. Especially in utero placental hemorrhage due to measures taken at birth is required.

### **Lactation**

Heparin does not appear in breast milk. Therefore, it is possible to breastfeed during treatment with heparin.

### **Reproductive ability/fertility**

There is not enough study.

### **4.7. Effects on ability to drive and use machines**

None stated.

### **4.8. Undesirable effects**

Undesirable events are listed by system organ class and frequency with the following approach: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); unknown (cannot be estimated from the available data)

The most frequently reported adverse events; hemorrhage, reversible increases in liver enzymes, reversible thrombocytopenia and various skin reactions. Isolated reports have been reported widespread allergic reactions, skin necrosis and priapism.

### **Blood and lymphatic system disorders**

Rare: Thrombocytopenia has been observed occasionally. (See Section 4.4)

Two types of heparin-induced thrombocytopenia have been defined:

Type I is frequent, mild (usually  $>50 \times 10^9/L$ ) and transient, occurring within 1-5 days of heparin administration.

Type II is less frequent but often associated with severe thrombocytopenia (usually  $<50 \times 10^9/L$ ). It is immunemediated and occurs after a week or more (earlier in patients previously exposed to heparin). It is associated with the production of a platelet-aggregating antibody and thromboembolic complications which may precede the onset of thrombocytopenia. Heparin should be discontinued immediately.



## Module 1 Administrative Information

### Module 1.3 Product Information

#### Module 1.3.1 SPC, PIL, Ampoule Print Form, The printing on the box

In some cases, thrombocytopenia type II is accompanied by venous or arterial thrombosis.

#### **Immune system disorders**

Rare: Any type and severe allergic reactions with various hypersensitivity symptoms (urticaria, conjunctivitis, rhinitis, asthma, cyanosis, tachypnoea, feeling of oppression, fever, chills)  
Anaphylactic reaction and anaphylactic shock  
Angioneurotic oedema

#### **Metabolism and nutrition disorders**

Rare: Heparin products can cause hypoaldosteronism which may result in an increase in plasma potassium. Rarely, clinically significant hyperkalemia may occur particularly in patients with chronic renal failure and diabetes mellitus (see Warnings and Precautions).

#### **Vascular disorders**

Rare: Haemorrhage. More may occur with high doses of heparin. (see also Special Warnings and Precautions and Overdosage Information).  
Very rare cases of epidural or spinal hematoma, prophylaxis of spinal or epidural anesthesia or spinal puncture implementation in patients receiving heparin has been reported.

#### **Hepatobiliary disorders**

Rare: Elevated transaminases, gamma-GT, LDH and lipase levels  
They usually resolve after discontinuation of heparin.

#### **Skin and subcutaneous tissue disorders**

Common: Rash (erythematous and various types rash such as maculopapular), urticaria, pruritus

Rare: Skin necrosis. Treatment should be stopped immediately if skin necrosis occurs.

#### **Musculoskeletal and connective tissue disorders**

Uncommon: There is some evidence that prolonged dosing with heparin (ie. over many months) may cause alopecia and osteoporosis. Significant bone demineralisation has been reported in women taking more than 10,000 I.U. per day of heparin for at least 6 months.

**Reproductive system and breast disorders**

Very rare: Priapism has been reported.

**General disorders and administration site conditions**

Very common: Injection site reactions; When injected into the skin under local irritation may occur.

**Reporting of the side effects**

If you get any side effects not listed in this leaflet, talk to your doctor or pharmacist.

You can also report side effects directly to your doctor or pharmacist. You can also report side effects directly to your country's related health authority. By reporting side effects, you can help provide more information on the safety of this medicine

**4.9. Overdose**

A potential hazard of heparin therapy is haemorrhage, but this is usually due to overdosage and the risk is minimised by strict laboratory control. Slight haemorrhage can usually be treated by withdrawing the drug. If bleeding is more severe, clotting time and platelet count should be determined. Prolonged clotting time will indicate the presence of an excessive anticoagulant effect requiring neutralisation by intravenous protamine sulphate, at a dosage of 1 mg for every 100 I.U. of heparin to be neutralised. The bolus dose of protamine sulphate should be given slowly over about 10 minutes and not exceed 50 mg. If more than 15 minutes have elapsed since the injection of heparin, lower doses of protamine will be necessary.

**5. PHARMACOLOGICAL PROPERTIES****5.1. Pharmacodynamic properties**

Pharmacotherapeutic group: Antithrombotic agents / heparin group

ATC code: B01AB01

Heparin is an anticoagulant agents at various stages of the effect of the normal coagulation system. It inhibits in vivo an in vitro reactions leading to blood clotting and the formation of fibrin clots. Antithrombin III (heparin co-factor) in plasma with heparin in small therapeutic dose combination inactivates activated factor X and inhibits trombosis by preventing the conversion of prothrombin to thrombin. This dose-dependently increased rate of inhibition. A greater amount of heparin, by in activating thrombin and preventing the conversion of fibrinogen to fibrin, may prevent further clot formation in the event of the occurrence of active thrombosis. Heparin also prevents the formation of a fibrin clot resistant by inhibiting the activation fibrin-stabilizing factor.

SELOPARIN 25000 IU/ 5 ml Solution for Injection

The heparin does not have fibrinolytic activity and therefore can not resolve the existing clot.

Bleeding time is usually unaffected by heparin. Clotting time is prolonged full therapeutic doses of heparin, unaffected at low doses.

## **5.2. Pharmacokinetic properties**

### **General features**

Immediately after intravenous injection, and 20-30 minutes after subcutaneous injection shows the anticoagulant effect.

#### Absorption:

Heparin does not indicate the normal absorption in oral, rectal, intramuscular and subcutaneous administration. Subcutaneous administration form acceptable prophylactic use only.

#### Distribution:

Heparin does not cross the placenta and breast milk.

#### Metabolism:

Heparin is metabolized in the liver. A fraction of heparin is neutralized with various factors such as platelet factor IV, plasma proteins (including fibrinogen), the reticuloendothelial system. Metabolites are inactive.

#### Elimination:

Plasma half-life of heparin varies widely between individuals depending on the dose and type of use and is considered as an average of 90 minutes.

Heparin is eliminated by the kidneys and is excreted unchanged a small amount with urine.

### **Characteristic features of the patients**

Elimination half-life time may vary depending on the presence of obesity, renal failure, malignancy, pulmonary embolism, liver dysfunction and infections.

The same amount of patients over 60 years of age dose compared to patients under 60 years old may have higher plasma levels of heparin and longer activated partial thromboplastin time.

## **5.3. Preclinical safety data**

There are no preclinical data of relevance to the prescriber which are additional to those already included in other sections.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1. List of excipients

Benzyl alcohol

Sodium chloride

Sodium hydroxide or hydrochloric acid

Water for injections

### 6.2. Incompatibilities

Any feasibility study was not carried out due to this product should not be mixed with other medicinal products.

Heparin is incompatible with many injectable preparations e.g. some antibiotics, opioid analgesics and antihistamines.

The following drugs are incompatible with heparin;

Alteplase, amikacin sulphate, amiodarone hydrochloride, ampicillin sodium, aprotinin, benzylpenicillin potassium or sodium, cefalotin sodium, chlorpromazine hydrochloride, ciprofloxacin lactate, cisatracurium besilate, cytarabine, dacarbazine, daunorubicin hydrochloride, diazepam, doxorubicin hydrochloride, droperidol, erythromycin lactobionate, gentamicin sulphate, haloperidol lactate, hyaluronidase, hydrocortisone sodium succinate, kanamycin sulphate, labetolol hydrochloride, meticillin sodium, methotrimeprazine, netilmicin sulphate, nicardipine hydrochloride, oxytetracycline hydrochloride, pethidine hydrochloride, polymyxin B sulphate, promethazine hydrochloride, streptomycin sulphate, tobramycin sulphate, triflupromazine hydrochloride, vancomycin hydrochloride and vinblastine sulphate.

Dobutamine hydrochloride and heparin should not be mixed or infused through the same intravenous line, as this causes precipitation.

Heparin and reteplase are incompatible when combined in solution. If reteplase and heparin are to be given through the same line this, together with any Y-lines, must be thoroughly flushed with a 0.9% saline or a 5% glucose solution prior to and following the reteplase injection.

### 6.3. Shelf life

24 months

### 6.4. Special precautions for storage

Please note that the application of the clear solution and the packaging is opened before.



Module 1 Administrative Information

Module 1.3 Product Information

Module 1.3.1 SPC, PIL, Ampoule Print Form, The printing on the box

Store at 25°C below room temperature and in the original packaging. Do not freeze.

The vial after once opening is available for a maximum of 28 days at 25°C.

Multidose vial is for single patient use.

**6.5. Nature and contents of container**

Bromobutyl stopper, red flip-off aluminum cap, type I colorless glass vial.

It is presented in boxes containing 1 vial.

**6.6. Special precautions for disposal and other handling**

Unused products or waste materials should be disposed of in accordance with the “Medical Waste Control Regulations” and “Packaging and Packaging Waste Control Regulations”.

**7. MARKETING AUTHORISATION HOLDER**

Haver Farma İlaç A.Ş.  
Akbaba Mah. Maraş Cad. No:52/2/1  
Beykoz / Istanbul / TURKEY  
Tel: (0216) 324 38 38  
Fax: (0216) 317 04 98

**8. MARKETING AUTHORISATION NUMBER(S)**

253/39

**9. DATE OF FIRST AUTHORISATION / RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 23.09.2013

Renewal of the authorisation: -

**10. DATE OF REVISION OF THE TEXT**